

first phase folds. Deformation by compaction is rejected because the overlying Lapa sandstone is not deformed.

The above features suggest that the sediments correspond to an esker. The deposit is part of the low-stand system tract of the 3<sup>rd</sup> order sequence S<sub>2</sub> described by Canuto et al. (2001). — ( *December 14, 2001* ) .

\* Supported by FAPESP, No. 97/13973-2.

\*\*E-mail: jrcanuto@usp.br

### THE THIRD ORDER SEQUENCE STRATIGRAPHY OF THE ITARARÉ SUBGROUP (LATE PALEOZOIC) ACROSS PARANÁ BASIN\*

JOSÉ R. CANUTO\*\*, PAULO R. SANTOS AND ANTONIO C. ROCHA-CAMPOS

Instituto de Geociências, USP, São Paulo, SP.

The Itararé Subgroup (late Paleozoic) on the eastern margin of the Paraná Basin is subdivided into seven 50-80m thick, type I, third order sequences that correspond to fining-coarsening upward cycles. Sequences are bounded at base by an erosional unconformity that at the eastern margin of the basin is marked either by glacially striated and/or glaciotectionised bedrock, or intraformational, glacially striated surfaces.

Our depositional model considers that each erosional surface, usually overlain by subglacial tillite, records an advance of a grounded glacier. Higher up the glaciogenic section includes glacioterrestrial, glaciomarine and marine beds (tillites, flow diamictites, proglacial rhythmites with dropstones, outwash sandstone, etc.), deposited during retreat of the ice margin. The sequences culminate with marine shale documenting short-lived post-glacial transgression, and shallow-marine progradational fluvial-deltaic sandstone, associated to glacioisostatic rebound that followed retreat of the glacial margin.

This type of sequence was earlier identified in outcrop and is now recognized in the subsurface of the Paraná Basin, along an east-west cross-section through wells 1-PT-1-PR, 1-RO-1-PR, 2-NA-1-PR and AMst-1-MT in Brazil, and Asunción 2 in Paraguay. Correlation of sequences in subsurface and outcrop was established lithostratigraphically from top to bottom, with reference to a datum represented by a widespread marine marker bed intercalated in the Rio Bonito Formation. Absence of basal S1 sequence in well 1-RO-1-PR may be related to its location on a basement high. Only the uppermost part of S1 is recorded in well AMst-1-MT due to the intercalation of a

thick diabase sill (Cretaceous) between Itararé beds and the underlying Ponta Grossa Formation.

With exception of the two cases, the subsurface section shows a full sequential correspondence. Most sequences, however, thicken from SE-NW, almost doubling values found on the eastern margin of the Paraná Basin. This suggests higher subsidence rate towards west, that compensated the positive tectonic behavior of the adjacent Asunción arch and created adequate space to accommodate sediments supplied from the eastern flank of the Asunción arch. — ( *December 14, 2001* ) .

\* Supported by FAPESP 97/13973-2.

\*\*E-mail: jrcanuto@usp.br

### EXTRUSION-RELATED EXTENSIONAL SIGNATURES UNDER HIGH-PRESSURE NAPPE COMPRESSIONAL REGIME\*

CARLOS H. G. CARVALHO AND MARIO C. CAMPOS-NETO  
Instituto de Geociências, USP, São Paulo, SP.

Presented by ANTONIO C. ROCHA-CAMPOS

The Neoproterozoic nappe system exposed south of the São Francisco Craton underwent a minimum of 200 km of near-horizontal northeastward displacement. This nappe system comprises the south-southeast sector of Brasília Belt and piles up three tectonic settings (from top to bottom): the roots of a neoproterozoic magmatic arc (*Socorro-Guaxupé Nappe*); high-pressure metasedimentary nappes (*Três Pontas-Varginha, Carmo da Cachoeira, Aiuruoca-Andrelândia, Pouso Alto and Carvalhos klippe*); parautochthonous medium-pressure nappes and duplexes.

In the southeast of Aiuruoca-Andrelândia Nappe, a brittle-ductile sinistral shear zone (*Liberdade Shear Zone*) limits the southern Alagoa Migmatites and may represent a lateral ramp for the eastern Lima Duarte Nappe. Close to Alagoa (MG), this shear zone turns E-W, breaking the nappe hind portion, in a southward normal-ductile movement.

North of this shear zone, the garnet-sillimanite bearing metasedimentary sequence lays over a gnaissic-diatexitic basement and is highly sliced by strata-like tourmaline-bearing granites. In the metasediments, main foliation S<sub>2</sub> is transposed by a normal-fault related thick protomylonitic S<sub>2</sub>' foliation. In the gnaisses, southward transport is marked by shear-sense indicators such as S-C foliations and quartz sigmoids. A low-temperature, late brittle thrust system, overlaps the extensional structures.

Heading north, the end of this structural-